

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : A61K 31/62, 31/54, 31/44 A61K 31/45, 31/40, 31/35	A1	(11) International Publication Number: WO 91/04031 (43) International Publication Date: 4 April 1991 (04.04.91)
(21) International Application Number: PCT/US90/04950 (22) International Filing Date: 5 September 1990 (05.09.90) (30) Priority data: 408,394 18 September 1989 (18.09.89) US (71) Applicant: E.I. DU PONT DE NEMOURS AND COMPANY [US/US]; 1007 Market Street, Wilmington, DE 19898 (US). (72) Inventors: ADDICKS, William, Joseph ; 732 Lora Lane, Hockessin, DE 19707 (US). MOLLICA, Joseph, A. ; 320 Centennial Circle, Greenville, DE 19807 (US). SLATKO, Gary, Howard ; 20 Carriage Pass, Chadds Fords, PA 19317 (US).		(74) Agents: HORN, Margaret, A. et al.; E.I. du Pont de Nemours and Company, Legal Department, 1007 Market Street, Wilmington, DE 19898 (US). (81) Designated States: AT (European patent), AU, BE (European patent), CA, CH (European patent), DE (European patent)*, DK (European patent), ES (European patent), FI, FR (European patent), GB (European patent), HU, IT (European patent), JP, KR, LU (European patent), NL (European patent), NO, SE (European patent), SU. Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: ORAL ANTICOAGULANT/PLATELET INHIBITOR LOW DOSE FORMULATION (57) Abstract There are disclosed combination products useful for the treatment and/or prevention of a first or recurrent myocardial infarction or a first or recurrent stroke. Said combination products comprise low doses of an oral anticoagulant, preferably warfarin and platelet inhibiting agents, including non-steroidal antiinflammatory agents and preferably aspirin.		

DESIGNATIONS OF "DE"

Until further notice, any designation of "DE" in any international application whose international filing date is prior to October 3, 1990, shall have effect in the territory of the Federal Republic of Germany with the exception of the territory of the former German Democratic Republic.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MC	Monaco
AU	Australia	FI	Finland	MG	Madagascar
BB	Barbados	FR	France	ML	Mali
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GR	Greece	NL	Netherlands
BJ	Benin	HU	Hungary	NO	Norway
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	SD	Sudan
CF	Central African Republic	KP	Democratic People's Republic of Korea	SE	Sweden
CG	Congo	KR	Republic of Korea	SN	Senegal
CH	Switzerland	LI	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
DE	Germany, Federal Republic of	LU	Luxembourg	TG	Togo
DK	Denmark			US	United States of America

TITLEORAL ANTICOAGULANT/PLATELET INHIBITORLOW DOSE FORMULATION5 Background of the InventionField of Invention

 This invention relates to combination products comprising low doses of oral anticoagulants, such as warfarin, and low doses of inhibitors of platelet
10 function (e.g., aggregation, adhesion) such as non-steroidal antiinflammatory agents, and preferably aspirin, in a single dosage form. Further, this invention relates to methods of using these low dose combination products for the prevention and/or treatment
15 of first or recurrent myocardial infarction or the prevention and/or treatment of first or recurrent stroke.

Background Information

 Recent advances in the understanding of the
20 pathogenic factors leading to the acute coronary ischemic syndromes of unstable angina, myocardial infarction and ischemic sudden death, and acute cerebrovascular ischemic syndromes like transient ischemic attacks and stroke have demonstrated the
25 individual importance of two compounds in particular, acetyl salicylic acid, hereinafter referred to as aspirin or ASA, and warfarin, in the prevention and/or treatment of these syndromes. Recently the value of antithrombotic therapy with high-dose aspirin in
30 unstable angina has been conclusively demonstrated by two randomized, placebo-controlled, double-blind trials. See generally Lewis, et al. "Protective Effects of Aspirin Against Acute Myocardial Infarction and Death in Men with Aspirin." Results of a Veterans Cooperative

Study, New England Journal of Medicine, 309: 396 (1983) and Cairns, et al.: "Aspirin, Sulfinpyrazone or Both in Unstable Angina: Results of a Canadian Multicenter Trial". New England Journal of Medicine, 313: 1369, 5 (1985). Also, the recently pooled results of several studies suggest that long-term high-dose oral anticoagulant therapy may reduce the rate of recurrence of myocardial infarction by about 20%. [Fuster, et. al, (1988) Perspective, "Insights into the Pathogenesis of 10 Acute Ischemic Syndromes", 77, No. 6, pp 1213-1220].

Since currently available pharmaceutical products have an isolated inhibitory effect on either platelet function or thrombus formation, the use of a combination of agents which would effect both platelet 15 aggregation/adhesion and thrombus formation simultaneously, provides a potential benefit of improved efficacy at lower dosages over the use of individual agents. Since aspirin may prevent ischemic cardiac events caused by coronary artery disease and oral 20 anticoagulants may protect against both ischemic cardiac events and resultant cerebral embolization from ventricular thrombi, the combination in small doses may provide the best overall protection. [Fuster, V., Halperin, J.L., (1989) The New England Journal of 25 Medicine, Feb. 9, 1989, pp 392-394.]

Traditionally, the simultaneous use of warfarin and aspirin at high doses has been relatively contraindicated. It has been a pervasive practice in the medical community to use these agents on an 30 either/or basis. This practice was largely due to medical literature reporting undesirable clinical and pharmacologic interactions of the two drugs at high doses. Clinically, the propensity of aspirin, at high doses, to cause gastric mucosal erosion/ulceration, when

dosed with an oral anticoagulant (warfarin) has led to a high reported incidence of exaggerated gastrointestinal (GI) bleeding with the high dose combination. Pharmacologically, aspirin at high doses also acts synergistically with warfarin to elevate the prothrombin time assay level for a given dose of warfarin. In light of recent advances in the study of the acute coronary syndromes and the recognition of the individual benefits that aspirin and warfarin provide in treating and/or preventing these syndromes, there is a need for a combination product wherein an oral anticoagulant such as warfarin and an antiplatelet agent, such as aspirin or a non-aspirin-non-steroidal antiinflammatory agent are present in a low dose ratio. The combination permits the use of doses below those currently accepted as "therapeutic" in the medical literature, in other words, doses at which the beneficial effects of the two agents are favored over the dose related side effects associated with simultaneous administration of currently accepted, high doses of the two agents.

Summary of the Invention

There are provided pharmaceutical compositions comprising an oral anticoagulant such as warfarin and a platelet inhibitor such as a non-steroidal antiinflammatory drug (NSAID) and more particularly a composition wherein the non-steroidal antiinflammatory drug is aspirin. The pharmaceutical compositions of this invention are combination products with low doses of each drug component. Further provided are pharmaceutical compositions comprising low doses of warfarin and a platelet inhibiting drug such as ticlopidine or other agents such as thromboxane receptor

antagonists or thromboxane synthetase inhibitor compounds. Still further provided is a method of using the pharmaceutical compositions of this invention for preventing and/or treating a first or recurrent myocardial infarction or first or recurrent stroke, in a mammal.

Detailed Description

The present invention is a pharmaceutical combination composition comprising warfarin present in an amount to be delivered of about 0.5 mg-10 mg and aspirin present in an amount to be delivered of about 10-500 mg. "Warfarin" as used herein means crystalline warfarin as well as amorphous sodium warfarin and derivatives thereof. Preferred compositions comprise warfarin present in an amount to be delivered of about 1-7.5 mg and aspirin present in an amount to be delivered of about 20-325 mg. The most preferred embodiments of the present invention comprise warfarin present in an amount to be delivered of about 1-5 mg and aspirin present in an amount to be delivered of about 40-162.5 mg.

Although aspirin is the preferred platelet inhibitory agent of the present invention, other non-aspirin, non-steroidal antiinflammatory agents that inhibit platelet function such as ibuprofen, naproxen, sulindac, indomethacin, mefenamate, droxicam, diclofenac, sulfinpyrazone and piroxicam, in combination with warfarin, are within the scope of this invention. Of the non-aspirin, non-steroidal antiinflammatory drugs (NSAIDS), a preferred embodiment of the present invention comprises warfarin and piroxicam, commercially available from Pfizer, as Feldene®, since piroxicam exerts an antiplatelet effect when dosed once daily. Preferably, a combination of piroxicam and warfarin

comprises warfarin present in an amount to be delivered of about 0.5 mg-10 mg and piroxicam present in an amount to be delivered of about 5-50 mg. More preferably the combination would contain warfarin present in an amount to be delivered of about 1-7.5 mg and piroxicam present in an amount to be delivered of about 10-40 mg. The most preferred embodiment comprises warfarin present in an amount to be delivered of about 1-5 mg and piroxicam present in an amount to be delivered of about 10-20 mg.

Other embodiments of this invention include the following NSAIDs present in the amount shown along with warfarin in the amount shown.

	Indomethacin	Warfarin
	20-100 mg	0.5-10 mg
15	Naproxen	Warfarin
	20-100 mg	0.5-10 mg
	Droxicam	Warfarin
	20-100 mg	0.5-10 mg
	Diclofenac	Warfarin
20	20-100 mg	0.5-10 mg
	Sulfinpyrazone	Warfarin
	20-100 mg	0.5-10 mg

More preferably the NSAIDs listed above are present in an amount of about 25-90 mg and warfarin is present in an amount of about 1-7.5 mg. The most preferred embodiment of non-aspirin-NSAIDs and warfarin comprises the NSAIDs listed above present in an amount of about 25-75 mg and warfarin present in an amount of about 1-5 mg.

Another embodiment of this invention includes other platelet inhibiting or thromboxane inhibiting compounds such as ticlopidine, in combination with warfarin. Platelet inhibitors such as ticlopidine are not

associated with the GI irritation that aspirin and other non-steroidal antiinflammatory agents may exhibit and therefore may be a more acceptable dosage form in individuals with a history of GI problems such as ulceration. Furthermore, such a combination would be useful in a patient having a sensitivity to aspirin or other non-steroidal antiinflammatory agents. Preferably a combination of ticlopidine and warfarin comprises warfarin present in an amount to be delivered of about 0.5-10 mg and ticlopidine present in an amount to be delivered of about 100-750 mg. More preferably, the combination comprises warfarin present in an amount to be delivered of about 1-7.5 mg and ticlopidine present in an amount to be delivered of about 200-600 mg. The most preferred embodiment comprises warfarin present in an amount to be delivered of about 1-5 mg and ticlopidine is present in an amount to be delivered of about 250-500 mg.

Other platelet aggregation or adhesion inhibitory agents such as thromboxane-A₂-receptor antagonists and thromboxane-A₂-synthetase inhibitors are also useful in the present invention.

The combination products of the present invention can be in any dosage form known to those skilled in the art, such as tablets either single or multi-layered, capsules, caplets, liquids, or suppositories. The preferred dosage forms are tablets, capsules and caplets. These preferred dosage forms can be made by methods known to those skilled in the art and described in Remington's Pharmaceutical Sciences, (1985), 17th Edition, Osol, a standard reference in the field.

Due to the nature of the preferred compounds of this invention, warfarin and aspirin, the potential exists for a chemical interaction between warfarin and

aspirin formulated in a single dosage form. This potential is evidenced by results of compatibility experiments in which aspirin and sodium warfarin were intimately mixed in the ratio of 1 part sodium warfarin to 80 parts aspirin and stored at room temperature, 60°C, or under 600 foot-candles of light. Results recorded as average % recovery (\pm s.d.) are shown in Table I.

10

Table I

Aspirin And Sodium Warfarin Solid
State Stability After One Week

15	Condition	Aspirin	Aspirin	Sodium
		Recovery	Recovery	Warfarin
20		Where	Where	Where
		Aspirin Is	Aspirin Is Mixed* With	Sodium Warfarin Is Mixed*
25		<u>Stored Alone</u>	<u>Sodium Warfarin</u>	<u>With Aspirin</u>
	room temperature (22°C)	101.1 (0.5)	100.5 (0.5)	99.6 (1.8)
	60°C	99.8 (1.6)	76.6 (2.9)	72.8 (3.31)
	600 F-C	100.9 (1.0)	94.1 (2.7)	88.8 (9.5)

*mixed at a ratio of 1 part sodium warfarin to 80 parts aspirin

30

For this reason, the preferred dosage forms of this invention are formulated such that the physical contact between the warfarin and the aspirin is minimized. In order to minimize the contact between the warfarin and aspirin components, one embodiment of this invention provides for a combination product wherein the aspirin component is enteric coated. By enteric coating the

35

aspirin, it is possible not only to minimize the contact between the two active ingredients, but also, it is possible to control the release of the aspirin in the GI tract such that the aspirin is not released in the stomach but rather is released in the intestines. Such a formulation will further reduce the risk of GI side effects associated with aspirin since not only will the aspirin be present in a low dose, but also, it will be enteric coated thereby releasing lower in the GI tract.

Another embodiment of this invention provides for a combination product wherein the aspirin is coated with a sustained-release material which effects a sustained release throughout the gastrointestinal tract and also serves to minimize physical contact between aspirin and warfarin. Furthermore, the sustained release aspirin can be additionally enteric coated such that aspirin release occurs only in the intestine.

Still another approach would involve the formulation of a combination product in which the aspirin component is coated with a sustained and/or enteric release polymer, and the warfarin particles or granules are also coated with a polymer such as a low-viscosity grade of hydroxypropyl methylcellulose (HPMC) or other appropriate materials as known in the art in order to further separate the active components. The warfarin can alternatively be dry granulated with appropriate excipients, and the resulting granules are then coated with HPMC. The coating of the warfarin component serves to form an additional barrier to interaction with aspirin.

Dosage forms of the present invention wherein the aspirin component is enteric coated can be in the form of tablets such that the enteric coated aspirin and the warfarin are blended together and then compressed into a

tablet or such that the enteric coated aspirin is compressed into one tablet layer and the warfarin is compressed into an additional layer. Optionally, in order to further separate the two layers, one or more
5 placebo layers may be present such that the placebo layer is between the warfarin layer and aspirin layer.

In addition, dosage forms of the present invention can be in the form of capsules wherein the aspirin component is compressed into a tablet or preferably the
10 aspirin component is in the form of a plurality of microtablets, particles, granules or non-pareils, which are then enteric coated. These enteric coated microtablets, particles, granules or non-pareils are then placed into a capsule or compressed into a capsule
15 along with the warfarin granulation. Conversely, the warfarin component can be made into a tablet or microtablets and placed into a capsule along with a powder blend consisting of enteric or sustained release aspirin present as coated particles, granules or non-
20 pareils.

Further embodiments of this invention comprise a sustained release dosage form in which aspirin is formulated in such a way as to be delivered over 3-12 hours. The aspirin can alternatively be formulated to
25 effect a 3-12 hour delivery in the intestine only.

The formulations of the present invention may comprise, in addition to active ingredients, appropriate excipients, including but not limited to: lubricants such as magnesium stearate, stearic acid, sodium stearyl
30 fumarate, and hydrogenated vegetable oil; diluents such as lactose, microcrystalline cellulose, dextrose, starch and dicalcium phosphate; a glidant such as colloidal silicon dioxide; and disintegrates such as sodium starch glycolate and croscarmellose sodium. The use of such

excipients is known to those skilled in the art of pharmaceutical formulations.

The following examples describe various formulations of warfarin and aspirin combination products. These formulations are merely exemplary, it being understood that those skilled in the art can make obvious modifications thereof.

I. Immediate Aspirin Release Within the Intestine

10 1. Enteric-Coated

Aspirin/Sodium Warfarin

The following strategies can be utilized to achieve a product yielding the delivery of drug directly to the intestine whereupon it is immediately released, as well as a physical separation of sodium warfarin and aspirin.

A. Enteric Coated Aspirin Blended
with Sodium Warfarin

Aspirin particles, granules, or non-pareils are enteric coated with either a solvent-based or water-based enteric coating polymer, using a fluidized-bed process, a pan-coating process, a solids processor, or other methods known in the art. Examples of enteric coatings which can be used are cellulose acetate phthalate (Aquateric®, FMC Corp.) and various methacrylic acid copolymers (Eudragit®L30D, Eudragit®L and/or Eudragit®S, available from Röhm Pharma.), and other enteric coat materials as known in the art. Eudragit®L30D may be used alone or in combination with Eudragit®NE 30D (a methacrylate ester copolymer available from Röhm Pharma). For the cases in which the coating process is carried out using an aqueous-based solvent, a thin subcoat of hydroxypropyl methylcellulose may be used. The coating layer serves to minimize contact between the aspirin and sodium warfarin within

the dosage form, in addition to effecting aspirin delivery to the intestine. These coated particles, granules, or non-pareils are then blended with sodium warfarin and appropriate excipients, and this mixture is subsequently either tableted or encapsulated. Following is an example of a formulation of this type:

Example 1

Enteric Coated Aspirin
Blended With Sodium Warfarin

10		<u>Amount (mg) per</u>
	<u>Component</u>	<u>Dosage Unit</u>
	Aspirin coated with 9-25% Eudragit® L 30D	549.45-666.66 (500 mg active)
15	Sodium Warfarin	1.0
	Microcrystalline Cellulose	82.74-199.95
20	Sodium Starch Glycolate	32.0
	Colloidal Silica	1.6
	Sodium Stearyl Fumarate	16.0
25	<hr/>	
	To Total 800.0 mg	

Prior to weighing, pass all ingredients through an appropriate screen as needed. Weigh all ingredients and mix the sodium warfarin with the colloidal silica, sodium starch glycolate and one-half of the microcrystalline cellulose for about 15 minutes in a twin-shell blender. Add one-half of the sodium stearyl fumarate and mix for about 3 minutes. Slug this blend

using 1/2 inch flat-face punches and mill. Add the aspirin to the remaining microcrystalline cellulose and mix for about 15 minutes. Add this blend to the sodium warfarin granulation and mix for about 7 minutes. Add the remaining sodium stearyl fumarate, mix for about 3 minutes, and compress into tablets. It should be noted that the microcrystalline cellulose in this formulation is adjusted such that the total dosage form weight is 800.0 mg.

10

Example 2Enteric Coated Aspirin
Blended With Sodium Warfarin

15	<u>Component</u>	<u>Amount (mg) per</u> <u>Dosage Unit</u>
	Aspirin coated with 9.4% Eudragit®L30D (2% hydroxy- propyl methylcellulose subcoat)	180.59 (160 mg active)
20	Starch	100.16
	Sodium Warfarin	1.0
25	Microcrystalline Cellulose	196.0
	Sodium Starch Glycolate	20.0
	Colloidal Silica	1.0
30	Magnesium Stearate	1.25

To Total 500.0 mg

Prior to weighing, pass all ingredients through an appropriate screen as needed. Weigh all ingredients and mix the sodium warfarin with the colloidal silica, starch, sodium starch glycolate and one-half of the microcrystalline cellulose for about 15 minutes in a twin-shell blender. Add one-half the magnesium stearate and blend for about 3 minutes. Slug this blend using 1/2 inch flat-face punches and mill. Add the aspirin to the remaining microcrystalline cellulose and blend for about 15 minutes. Add this blend to the sodium warfarin granulation, and blend for about 7 minutes. Add the remaining magnesium stearate, mix for about 3 minutes, and compress into tablets or alternatively, place into capsules.

Using this formula and manufacturing directions, 7/16 inch tablets were produced, having a hardness of 8-10 Strong-Cobb Units (SCU). Dissolution tests were performed for both drugs in accordance with the dissolution tests for enteric coated aspirin tablets and sodium warfarin tablets as described in the United States Pharmacopeia/National Formula (USP/NF) XVII. For aspirin, dissolution was carried out by placing tablets in baskets rotating at 100 rpm. The dissolution media consisted of 750 ml of 0.1N HCl (37°C) for the first two hours of the test, followed by the addition of 250 ml of 0.2 N Tribasic sodium phosphate for a resulting pH of 6.8 during the remainder of the test. All analysis was by reversed-phase high performance liquid chromatography (HPLC). Results of aspirin dissolution are shown in Table II.

Table II

Aspirin Dissolution From Tablets Manufactured
In Accordance With Example 2

	<u>Time (min)</u>	<u>pH</u>	<u>Average (n=3) % Dissolved</u>
5	60	1.0	0.7
	120	1.0	1.6
	180	6.8	100.3
	210	6.8	100.7

- 10 These tablets pass the criteria outlined in USP/NF XVII for enteric release aspirin tablets (less than 10% aspirin released after 120 minutes at pH 1.0, greater than 80% released after 90 minutes at pH 6.8).

- 15 Sodium warfarin dissolution was assessed by placing tablets in 900 ml of pH 7.5-phosphate buffer maintained at 37°C, using a paddle at 50 rpm. An average of three tablets yielded 91.3% sodium warfarin release in 30 minutes, which passes the specifications as set forth in USP/NF XVII (not less than 80% release in 30 minutes).

20

B. Enteric Coated Aspirin Compressed in a
Multilayer Tablet

- 25 This approach serves to greatly minimize the drugs' area of contact. Aspirin particles, granules, or non-pareils are first enteric coated as specified in I-1A. The enteric coated aspirin is blended with appropriate excipients, and the sodium warfarin is dry granulated with the same excipients. The aspirin blend is then compressed, and the sodium warfarin layer is
- 30 subsequently compressed onto this layer. These two drug layers may optionally be further separated by an additional placebo layer. Following is an example of a formulation of this type:

Example 3Enteric Coated Aspirin/Sodium
Warfarin Into A Multilayer Tablet

5	<u>Component</u>	<u>Amount (mg) per</u>
		<u>Dosage Unit</u>
	Aspirin coated with 9-25% Eudragit®L30D	43.96-53.33 (40.0 mg active)
10	Sodium Warfarin	5.0
	Lactose	324.62-333.99
	Sodium Stearyl Fumarate	8.0
15	Tapioca Starch, Pregelatinized	9.05
<hr/>		
20	To Total 400.0 mg	

Prior to weighing, pass all ingredients through an appropriate screen as needed. Weigh all ingredients and combine the sodium warfarin with one-half of the tapioca starch and lactose. Mix in a twin-shell blender for about 15 minutes. Add one-fourth of the sodium stearyl fumarate and mix for about 3 minutes. Slug this blend using 1/2 inch flat-face punches and mill. Add one-fourth of the sodium stearyl fumarate and mix for about 3 minutes. Mix the aspirin with the remainder of the lactose and tapioca starch for about 10 minutes in a twin-shell blender. Add the remaining sodium stearyl fumarate and mix for about 3 minutes. Using a multilayer tablet machine such as the Manesty Layer

Press available from Thomas Engineering, compress the aspirin blend, followed by the sodium warfarin granulation, to produce a bilayer tablet. It should be noted that the lactose in this formulation is adjusted
5 such that the total dosage form weight is 400.0 mg.

C. Enteric Coated Aspirin Tablet Within a Capsule

This formulation involves the manufacture of an aspirin tablet or aspirin microtablets which are then
10 enteric coated and placed into a capsule along with a sodium warfarin granulation. Following is an example of a formulation of this type:

Example 4Enteric Coated Aspirin Tablet/
Sodium Warfarin Blend Within A Capsule

5	<u>Component</u>	<u>Amount (mg) per Dosage Unit</u>
	Aspirin	40.0
10	Eudragit®L30D	24.0
	Sodium Warfarin	1.0
	Lactose	106.5
15	Microcrystalline Cellulose	39.0
	Dicalcium Phosphate	184.5
20	Croscarmellose Sodium	16.0
	Colloidal Silica	1.0
	Hydrogenated Vegetable Oil	12.0
25		

To Total 424.00 mg

Prior to weighing, pass all ingredients through an appropriate screen as needed. Weigh all ingredients and mix the aspirin with the lactose, microcrystalline cellulose and one-half of the colloidal silica and

5 croscarmellose sodium for about 15 minutes in a twin-shell blender. Add one-half of the hydrogenated vegetable oil and mix for about 5 minutes. Compress this blend into tablets, and coat these with Eudragit®L30D using an appropriate method known in the

10 art. Mix the sodium warfarin with the dicalcium phosphate and the remaining colloidal silica and croscarmellose sodium for about 15 minutes. Add one-fourth of the hydrogenated vegetable oil and mix for about 5 minutes. Slug this blend using 1/2 inch flat-

15 face punches and mill. Add the remaining hydrogenated vegetable oil and mix for about 5 minutes. Encapsulate this formulation with the use of a capsule filling machine such as the Zanasi AZ5, commercially available from Soteco USA, Inc., by placing 200 mg of the sodium

20 warfarin granulation and one aspirin tablet into each capsule.

Example 5Sodium Warfarin Tablet/Enteric
Coated Aspirin Blend Within A Capsule

5	<u>Component</u>	<u>Amount (mg) per Dosage Unit</u>
10	Aspirin coated with 9-25% Eudragit®L30D	82.42 - 100.0 (75 mg active)
	Sodium Warfarin	10.0
	Lactose	57.5-75.08
15	Microcrystalline Cellulose	20.0
	Sodium Starch glycolate	8.0
	Colloidal Silica	0.5
20	Sodium Stearyl Fumarate	4.0
<hr/>		
To Total 200.0 mg		

- 25 Prior to weighing, pass all ingredients through an appropriate screen as needed. Weigh all ingredients and mix the aspirin with one-half of the lactose, sodium starch glycolate and colloidal silica for about 15 minutes in a twin-shell blender. Add one-half of the
- 30 sodium stearyl fumarate and mix for about 5 minutes. Combine the sodium warfarin with the microcrystalline cellulose and the remaining lactose, sodium starch glycolate and colloidal silica and blend for about 15 minutes. Add one-fourth of the sodium stearyl fumarate

and blend for about 4 minutes. Slug this blend using
1/2 inch flat-face punches and mill. Add the remainder
of the sodium stearyl fumarate and blend for about 4
minutes. Compress the sodium warfarin granulation into
5 tablets. Encapsulate this formulation with the use of a
capsule filling machine such as the Zanasi AZ5, by
placing one tablet along with a sufficient amount of
aspirin granulation to equal one dosage unit into each
capsule. It should be noted that the amount of lactose
10 in this formulation is adjusted such that the total
dosage form weight is 200 mg.

II. Sustained Aspirin Release

1. Sustained Release Aspirin Particles/Sodium 15 Warfarin

These formulations involve the coating of aspirin
microtablets, particles, granules, or non-pareils with a
sustained-release material such as methacrylate ester
copolymers (Eudragit®NE30D, Eudragit®RS30D and/or
20 Eudragit®RL30D, Eudragit®RL and/or Eudragit®RS, all
available from Röhm Pharma), organic-based
ethylcellulose, aqueous-based ethylcellulose (Aquacoat®,
FMC Corp.; Surelease®, Colorcon), or other appropriate
material as known in the art to effect a sustained
25 release of 3-12 hours. These coated entities can be
additionally overcoated with a layer of low viscosity
hydroxypropylmethylcellulose in order to guard against
rupture of the sustaining layer upon compression. These
microtablets, particles, granules, or non-pareils can
30 optionally be additionally enteric coated with an
enteric release polymer such as Aquateric®,
Eudragit®L30D, or Eudragit®L and/or Eudragit®S to effect
a sustained release in the intestine only. Another
approach involves the use of a combination of polymers

(such as Eudragit®NE30D and Eudragit®L30D) in a single coating to produce sustained release in the intestine. These layers can be applied by fluidized bed, coacervation-phase separation, pancoating, solids processor or other appropriate methods as known in the art. The coated aspirin is then mixed with appropriate excipients, and this blend is mixed with a sodium warfarin granulation and either placed into a capsule or compressed into a tablet. Alternatively, the sustained release aspirin blend can be compressed into a tablet or microtablets, which optionally may be additionally coated with an enteric layer, and placed into a capsule along with the sodium warfarin granulation. These two mixtures can also be processed in such a way as to produce a multilayer tablet with the aspirin and sodium warfarin existing in separate, distinct layers. These alternatives yield products which release the aspirin in a sustained manner. Additionally, contact between the aspirin and sodium warfarin within the product is minimized. The following is an example of a formulation of this type:

Example 6Sustained Release Aspirin/Sodium Warfarin Blend

	<u>Component</u>	<u>Amount (mg) per</u> <u>Dosage Unit</u>
5	Aspirin (coated with 3-15% Surelease® (ethylcellulose))	164.95-188.24 (160.0 mg active)
	Sodium Warfarin	5.0
10	Microcrystalline Cellulose	184.76-208.05
	Sodium Starch Glycolate	16.0
15	Colloidal Silica	1.0
	Sodium Stearyl Fumarate	8.0
20	<hr/>	
	To Total	403.00 mg

Prior to weighing, pass all ingredients through an appropriate screen as needed. Weigh all ingredients and

25 place the sustained release coated aspirin, one-half of the microcrystalline cellulose, colloidal silica and sodium starch glycolate into a twin-shell blender, and mix for about 15 minutes. Combine the sodium warfarin with the remaining microcrystalline cellulose, colloidal

30 silica, and sodium starch glycolate and mix for about 15 minutes. Add one-half of the sodium stearyl fumarate to the sodium warfarin blend and blend for about 3 minutes. Slug this blend using 1/2 inch flat-face punches and mill. Combine the aspirin blend with the sodium

- warfarin granulation and mix for about 10 minutes in a twin-shell blender. Add the remaining sodium stearyl fumarate, mix for about 3 minutes and either compress the blend into tablets or place into capsules. It should be noted that the microcrystalline cellulose in this formulation is adjusted such that the total dosage form weight is 403.0 mg.

Example 7

- 10 Sustained Release Aspirin
 Compressed Into A Tablet (Or
 Microtablets)/Sodium Warfarin In A Capsule
 Using the ingredients of Example 4, in like
 quantity, prior to weighing, pass all ingredients
15 through an appropriate screen as needed. Weigh all
 ingredients and place the sustained release coated
 aspirin and one-half of the microcrystalline cellulose,
 sodium starch glycolate, and colloidal silica into a
 twin-shell blender and mix for about 15 minutes. Add
20 one-half of the sodium stearyl fumarate and mix for
 about 5 minutes. Compress this mixture into a tablet or
 microtablets. Combine the sodium warfarin with the
 remaining microcrystalline cellulose, colloidal silica,
 and one-fourth of the sodium stearyl fumarate and mix
25 for about 15 minutes. Slug this blend using 1/2 inch
 flat-face punches and mill. Add the remaining sodium
 stearyl fumarate and mix for about 5 minutes. With the
 use of a capsule filling machine such as the Zanasi A25,
 encapsulate this formulation by placing an aspirin
30 tablet or microtablets and an amount of sodium warfarin
 granulation for one dosage unit into each capsule.

Example 8Sustained Release Aspirin/Sodium
Warfarin In A Multilayered Tablet

	<u>Component</u>	<u>Amount (mg) per Dosage Unit</u>
5	Aspirin (coated with 3-15% Surelease® (ethylcellulose))	164.95-188.24 (160.0 mg active)
	Sodium Warfarin	5.0
10	Microcrystalline Cellulose	369.26-392.55
	Sodium Starch Glycolate	24.0
15	Colloidal Silica	1.5
	Sodium Stearyl Fumarate	12.0
20	<hr/> To Total 600.00 mg	

- Prior to weighing, pass all ingredients through an appropriate screen as needed. Weigh all ingredients and
- 25 place the sustained release coated aspirin and one-half of the microcrystalline cellulose, sodium starch glycolate and colloidal silica into a twin-shell blender and mix for about 15 minutes. Add two-thirds of the sodium stearyl fumarate and mix for about 5 minutes.
- 30 Combine the sodium warfarin with the remaining microcrystalline cellulose, sodium starch glycolate and colloidal silica and mix for about 15 minutes. Add one-sixth of the sodium stearyl fumarate and mix for about 5 minutes. Slug this blend using 1/2 inch flat-face
- 35 punches, and mill. Add the remaining sodium stearyl

fumarate and mix for about 5 minutes. Using a multilayer tablet machine such as the Manesty Layer Press (available from Thomas Engineering), compress the aspirin blend, followed by the sodium warfarin granulation, to produce a bilayer tablet.

2. Sustained-Release Matrix Tablet/Sodium Warfarin Within a Capsule

In this formulation a sustained release matrix tablet containing aspirin is first manufactured by methods known in the art, and this tablet is placed into a capsule along with a sodium warfarin granulation. The sustained release matrix is produced by mixing aspirin with various types of hydroxypropyl methylcellulose (HPMC) (Methocel®K, Methocel®J, Methocel®E, all available from Dow Chemical), and hydroxypropyl ethylcellulose (Klucel®, Dow Chemical), either alone or in combination, along with appropriate excipients. When using Methocel®K or Methocel®J having a 2% viscosity greater than 800 cps either alone or in combination, the combined amount of Methocel®K and/or Methocel®J should be greater than 25.8% of the total dosage form weight. If the Methocel®K or Methocel®E or a combination thereof have a 2% viscosity of less than 800 cps, then the hydroxypropyl content is less than 9% of the total dosage form weight. Alternatively, the Methocel®K and/or Methocel®J can be mixed with a non-HPMC cellulose ether provided that the non-HPMC cellulose ether comprises greater than 30% of the entire polymer mixture.

The matrix tablet can additionally be coated with a low-viscosity polymer (e.g., hydroxypropyl methylcellulose) which works in concert with the polymer within the matrix to yield a sustained release.

In all cases drug release takes place over about a 3-12 hour period. The sustained release tablet is then placed into a capsule along with a sodium warfarin granulation. Alternatively, the aspirin tablet can be
5 coated by methods known in the art with an enteric coating material such as Eudragit®L30D, Eudragit®L, Eudragit®S, Aquateric®, or other materials known in the art. The net result of this option is to effect a sustained release in the intestine only.

10

Example 9

Sustained-Release Matrix

Tablets/Sodium Warfarin Within A Capsule

15	<u>Component</u>	<u>Amount (mg) per</u> <u>Dosage Unit</u>
	Aspirin	160.0
	Methocel®E4M Premium CR	109.95
20	Sodium Warfarin	2.0
	Lactose	128.07
25	Dicalcium Phosphate	125.32
	Colloidal Silica	2.34
	Sodium Stearyl Fumarate	12.32
30	<hr/>	
	To Total	540.0 mg

Prior to weighing, pass all ingredients through an appropriate screen as needed. Mix the aspirin with the lactose, Methocel® and one-half of the colloidal silica for about 15 minutes in a twin-shell blender. Add two-thirds of the sodium stearyl fumarate and mix for an additional 5 minutes. Compress this blend into tablets (135.77 mg each) such that three tablets yields aspirin sufficient for one dosage unit (160 mg total aspirin).

Mix the sodium warfarin with the dicalcium phosphate and remaining colloidal silica for about 15 minutes. Add one-sixth of the sodium stearyl fumarate and mix for about 5 minutes. Slug this blend using 1/2 inch flat-face punches and mill. Add the remaining sodium stearyl fumarate and mix for about an additional 5 minutes. Encapsulate this formulation by placing sodium warfarin granulation to equal one dosage unit and three aspirin tablets into each capsule with the use of a capsule filling machine such as the Zanasi AZ5.

The aspirin tablet component of this formulation was manufactured (8/32 inch tablets, hardness 10-12 SCU), and drug release was assessed. The dissolution apparatus and media used were the same as that described in Example 2 with the exception that 500 ml of 0.1 N HCl was used during the first 2 hours, followed by adjustment of the pH to 7.5 with the addition of 250 ml of tribasic sodium phosphate. Dissolution data are shown in Table III.

Table III

Aspirin Release From Matrix Tablets
Manufactured In Accordance With Example 9

5			Average
	Time		(n=3)
	<u>(hour)</u>	<u>pH</u>	<u>% Released</u>
	1	1.0	38.3
10	2	1.0	56.5
	3	7.5	75.3
	4	7.5	84.7
	5	7.5	95.2
	6	7.5	99.6
15			

By substituting non-aspirin NSAID agents, ticlopidine, thromboxane receptor antagonists or thromboxane synthetase inhibitors for aspirin in the above examples, other formulations within the scope of this invention can be prepared. Additionally, other oral anticoagulant agents such as coumarin derivatives (for example phenindione, bishydroxycoumarin (dicumarol) and phenprocoumon can be substituted for warfarin in the above examples, rendering formulations within the scope of this invention.

Utility

The potential for enhanced efficacy and an improved margin of safety by combining a low-dose of an oral anticoagulant, such as warfarin and a low-dose of platelet inhibitor agents, in general or non-steroidal anti-inflammatory agents, particularly aspirin, is supported by the previous clinical experience with the use of each (or a closely related compound) individually

at high doses in the prevention and treatment of acute coronary ischemic syndromes and ischemic cerebrovascular disease, as well as by certain suggestive evidence from their simultaneous administration.

5 Clinical studies using oral anticoagulants alone, including crystalline sodium warfarin, have provided evidence of their efficacy in the treatment or secondary prevention of coronary artery disease. Of three published, randomized, controlled trials of the
10 treatment of acute myocardial infarction, oral anticoagulation significantly reduced overall mortality and the frequency of reinfarction in one study. Of the four published large, randomized, controlled trials of oral anticoagulants in the secondary prevention of
15 myocardial infarction, three suggested a reduction in the incidence of reinfarction and early mortality. One additional study, the Warfarin Reinfarction Study, has also recently demonstrated a significant reduction in mortality, reinfarction, and stroke in people with a
20 previous myocardial infarction who were treated with warfarin as compared to those treated with placebo.

 The results of studies utilizing acetylsalicylic acid (ASA) alone in the prevention and treatment of coronary artery disease have also been promising. In
25 patients with unstable angina, ASA has been demonstrated to reduce the incidence of subsequent myocardial infarction and mortality in two large, randomized, double-blind, placebo-controlled clinical studies. In addition, ASA has been approved for use in the secondary
30 prevention of myocardial infarction, based on data from several trials which, when pooled, suggested a reduction in reinfarction and mortality. Finally, two recent studies evaluating ASA in the primary prevention of coronary artery disease have reported either a dramatic

or inconsequential benefit; ASA has not as yet been approved for this indication.

In addition to their utility in coronary artery disease, agents that inhibit platelet function such as ASA and ticlopidine have been shown to be effective in the prevention of stroke in people with cerebrovascular disease. Pooled data from nine randomized trials have provided overwhelming evidence of the efficacy of ASA alone in reducing the risk of completed stroke in people with transient ischemic attacks (TIAs). Recently, ticlopidine alone has also been demonstrated to have efficacy in treating TIAs.

The postulate that warfarin and ASA in combination will afford an efficacy advantage over either agent alone is based on recent angiography data and is supported by both pilot studies and anecdotal reports. Based on the direct visualization of ulcerated atherosclerotic plaques within the coronary arteries, Forrester et al. have proposed two distinct pathologic cycles of platelet-dependent and thrombus-dependent injury and repair in acute coronary ischemic syndromes which strongly suggest a potential benefit of the two agents when used in combination. This pathophysiologic pattern can be extrapolated to the cerebrovascular system, as well. Recent pilot data in patients with unstable angina revealed a benefit in patients receiving both heparin (an anticoagulant)/warfarin and ASA, when compared to those receiving either agent alone. Additionally, patients with prosthetic heart valves, receiving both agents simultaneously for the prevention of stroke, have been noted to have a remarkably low incidence of myocardial infarction.

There is a need for a therapeutic approach to the acute coronary ischemic syndromes and ischemic

cerebrovascular diseases which takes into account the diverse pathophysiologic makeup of such diseases and which includes therapeutic agents for ameliorating each of these pathophysiological components. A combination
5 product containing a low dose of warfarin and a low dose of a platelet-inhibiting agent such as a non-steroidal antiinflammatory agent, and particularly aspirin, can provide such an approach. In addition, by administering low doses of each, the incidence of GI and other side
10 effects associated with the concurrent administration of these drugs at higher doses is significantly reduced and the patient is offered a convenient single dosage form for the once-daily, lifelong administration of these two medications. It is generally accepted that increased
15 convenience to the patient results in an increase in compliance. Furthermore, a combination product, available by prescription only, will give the physician greater dose control than if the patient were to independently select from the various dosages and
20 formulations and purchase aspirin over-the-counter. The combination products of the present invention will also reduce the likelihood of patient confusion often associated with concurrent dosing of medication when it is not available in a combination product.

What is Claimed is:

1. A pharmaceutical composition comprising a suitable pharmaceutical carrier and a combination of active ingredients consisting essentially of an oral
5 anticoagulant and a platelet inhibitory agent, said active ingredients being present in a ratio which provides therapeutic effectiveness with limited dose related side effects.
2. A composition of claim 1 wherein the oral
10 anticoagulant is warfarin or a pharmaceutically acceptable salt thereof.
3. A composition of claim 1 wherein the platelet inhibitory agent is acetylsalicylic acid (ASA).
4. A composition of claim 1 wherein the oral
15 anticoagulant is warfarin or a pharmaceutically acceptable salt thereof and the platelet inhibitory agent is acetylsalicylic acid.
5. A composition of claim 4 wherein the warfarin is present in an amount to be delivered of about 0.5-10
20 mg and the acetylsalicylic acid is present in amount to be delivered of about 10-500 mg.
6. A composition of claim 5 wherein the warfarin is present in an amount to be delivered of about 1-7.5 mg and the acetylsalicylic acid is present in an amount
25 to be delivered of about 20-325 mg.
7. A composition of claim 6 wherein the warfarin is present in an amount to be delivered of about 1-5 mg and the acetylsalicylic acid is present in an amount to be delivered of about 40-162.5 mg.
- 30 8. A composition of claim 4 wherein the combination is in a dosage form which provides minimal physical contact between the warfarin and the acetylsalicylic acid.

9. A composition of claim 8 wherein to achieve the minimal contact, the acetylsalicylic acid component is enteric coated.

10. A composition of claim 8 wherein to achieve the minimal contact, the acetylsalicylic acid component is coated with sustained release material and optionally further enteric coated.

11. A composition of claim 1 wherein the platelet inhibitory agent is a non-aspirin-non-steroidal antiinflammatory agent selected from the group consisting of piroxicam, indomethacin, naproxene, droxicam, diclofenac and sulfinpyrazone.

12. A composition of claim 2 wherein the platelet inhibitory agent is a non-aspirin-non-steroidal antiinflammatory agent selected from the group consisting of piroxicam, indomethacin, naproxene, droxicam, diclofenac and sulfinpyrazone.

13. A composition of claim 12 wherein the non-aspirin-non-steroidal antiinflammatory agent is piroxicam.

14. A composition of claim 13 wherein the warfarin is present in an amount to be delivered of about 0.5-10 mg and the piroxicam is present in an amount to be delivered of about 5-50 mg.

15. A composition of claim 12 wherein the non-aspirin-non-steroidal antiinflammatory agent is indomethacin.

16. A composition of claim 15 wherein the warfarin is present in an amount to be delivered of about 0.5-10 mg and the indomethacin is present in an amount to be delivered of about 20-100 mg.

17. A composition of claim 12 wherein the non-aspirin-non-steroidal antiinflammatory agent is naproxene.

18. A composition of claim 17 wherein the warfarin is present in an amount to be delivered of about 0.5-10 mg and the naproxene is present in an amount to be delivered of about 20-100 mg.

5 19. A composition of claim 12 wherein the non-aspirin-non-steroidal antiinflammatory agent is droxicam.

10 20. A composition of claim 17 wherein the warfarin is present in an amount to be delivered of about 0.5-10 mg and the droxicam is present in an amount to be delivered of about 20-100 mg.

21. A composition of claim 12 wherein the non-aspirin-non-steroidal antiinflammatory agent is diclofenac.

15 22. A composition of claim 21 wherein the warfarin is present in an amount to be delivered of about 0.5-10 mg and the diclofenac is present in an amount to be delivered of about 20-100 mg.

20 23. A composition of claim 12 wherein the non-aspirin-non-steroidal antiinflammatory agent is sulfinpyrazone.

25 24. A composition of claim 23 wherein the warfarin is present in an amount to be delivered of about 0.5-10 mg and the sulfinpyrazone is present in an amount to be delivered of about 20-100 mg.

25. A composition of claim 1 wherein the platelet inhibitory agent is ticlopidine.

26. A composition of claim 2 wherein the platelet inhibitory agent is ticlopidine.

30 27. A composition of claim 26 wherein the warfarin is present in an amount to be delivered of about 0.5-10 mg and the ticlopidine is present in an amount to be delivered of about 100-750 mg.

28. A composition of claim 27 wherein the warfarin is present in an amount to be delivered of about 1-7.5 mg and the ticlopidine is present in an amount to be delivered of about 200-600.

5 29. A composition of claim 28 wherein the warfarin is present in an amount to be delivered of about 1-5 mg and the ticlopidine is present in an amount to be delivered of about 250-500 mg.

10 30. A composition of claim 1 in a tablet dosage form.

31. A composition of claim 1 in a capsule dosage form.

32. A composition of claim 1 in a caplet dosage form.

15 33. A composition of claim 1 in a suppository form.

34. A pharmaceutical composition comprising:

20 (a) about 40 to about 162.5 mg of active acetylsalicylic acid which is optionally coated with an appropriate enteric release polymer;

(b) about 1 to about 7.5 mg of warfarin or a pharmaceutically acceptable salt thereof which is optionally coated with an appropriate polymer to minimize contact between the active components of a) and
25 b); and

(c) a pharmaceutically acceptable carrier.

35. A pharmaceutical composition of claim 34 wherein the acetylsalicylic acid is coated with an enteric release polymer selected from the group
30 consisting of cellulose acetate phthalate or methacrylic acid copolymers.

36. A pharmaceutical composition of claim 34 wherein the warfarin is coated with a low-viscosity grade of hydroxypropyl methylcellulose.

37. A pharmaceutical composition of Claim 35 wherein the warfarin is coated with a low-viscosity grade of hydroxypropyl methylcellulose.

38. A pharmaceutical composition comprising:

- 5 (a) about 40 to about 162.5 mg of active acetylsalicylic acid which is optionally coated with an appropriate sustained release polymer;
- (b) about 1 to about 7.5 mg of warfarin which is optionally coated with an appropriate polymer to
- 10 minimize contact of the active components steps a) and b); and
- c) a pharmaceutically acceptable carrier.

39. A pharmaceutical composition of claim 38 wherein the acetylsalicylic acid is coated with an

15 appropriate sustained release polymer such that release occurs over about 3 to 12 hours.

40. A pharmaceutical composition of claim 39 wherein the sustained release polymer is selected from the group consisting of methacrylate ester copolymers,

20 organic-based ethylcellulose or aqueous-based ethylcellulose.

41. A pharmaceutical composition of claim 38 wherein the acetylsalicylic acid is additionally coated with an appropriate enteric release polymer.

25 42. A pharmaceutical composition of claim 41 wherein the enteric release polymer is methacrylic acid copolymer.

43. A method of preventing and/or treating a first or recurrent myocardial infarction or preventing and/or

30 treating a first or recurrent stroke in a mammal comprising administering to the mammal an effective amount of any of the compositions of claims 1 to 29 or 34 to 42.

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US90/04950

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ⁸ According to International Patent Classification (IPC) or to both National Classification and IPC IPC (5): A61K 31/62, A61K 31/54, A61K 31/44, A61K 31/45, A61K 31/40, A61K 31/35 U.S. CL.: 514/161, 226.5, 301, 404, 420, 457, 420, 822																							
II. FIELDS SEARCHED <div style="text-align: center; border-top: 1px solid black; border-bottom: 1px solid black;">Minimum Documentation Searched ⁷</div> <table style="width: 100%; border-collapse: collapse;"> <tr> <th style="width: 25%; text-align: left; border-bottom: 1px solid black;">Classification System</th> <th style="width: 75%; text-align: left; border-bottom: 1px solid black;">Classification Symbols</th> </tr> <tr> <td style="height: 40px; vertical-align: top; border-right: 1px solid black; border-bottom: 1px solid black;">U.S.</td> <td style="border-bottom: 1px solid black;">514/161, 226.5, 301, 404, 420, 457, 822</td> </tr> </table> <div style="text-align: center; border-top: 1px solid black; border-bottom: 1px solid black;">Documentation Searched other than Minimum Documentation to the extent that such Documents are included in the Fields Searched ⁸</div>			Classification System	Classification Symbols	U.S.	514/161, 226.5, 301, 404, 420, 457, 822																	
Classification System	Classification Symbols																						
U.S.	514/161, 226.5, 301, 404, 420, 457, 822																						
III. DOCUMENTS CONSIDERED TO BE RELEVANT ⁹ <table style="width: 100%; border-collapse: collapse;"> <tr> <th style="width: 10%; text-align: left; border-bottom: 1px solid black;">Category ⁹</th> <th style="width: 60%; text-align: left; border-bottom: 1px solid black;">Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²</th> <th style="width: 30%; text-align: left; border-bottom: 1px solid black;">Relevant to Claim No. ¹³</th> </tr> <tr> <td style="text-align: center; vertical-align: top; border-right: 1px solid black;">X</td> <td style="border-right: 1px solid black;">FR, 2265-273 (LAB SOOTHERAP) 29 March 1974 See the translation</td> <td style="vertical-align: top;">1-43</td> </tr> <tr> <td style="text-align: center; vertical-align: top; border-right: 1px solid black;">Y</td> <td style="border-right: 1px solid black;">FR, 2265-273 (LAB ZOOTHERAP) 29 March 1974 See the translation</td> <td style="vertical-align: top;">1-43</td> </tr> <tr> <td style="text-align: center; vertical-align: top; border-right: 1px solid black;">Y</td> <td style="border-right: 1px solid black;">US, A, 4,444,777 (FLEMING, JR) 24 April 1984 See column 3, 4, and 5</td> <td style="vertical-align: top;">1-43</td> </tr> <tr> <td style="text-align: center; vertical-align: top; border-right: 1px solid black;">Y</td> <td style="border-right: 1px solid black;">US, A, 4,537,894 (BLANCHARD) 27 August 1985 See column 4 lines 1-15.</td> <td style="vertical-align: top;">1-43</td> </tr> <tr> <td style="text-align: center; vertical-align: top; border-right: 1px solid black;">Y</td> <td style="border-right: 1px solid black;">FR, 2390-959 (BRUGNAUD) 01 January 1979 See the translation.</td> <td style="vertical-align: top;">1-43</td> </tr> <tr> <td style="text-align: center; vertical-align: top; border-right: 1px solid black;">Y</td> <td style="border-right: 1px solid black;">EP 0001-503 (NUNN) 18 April 1979 See page 2, lines 14-21</td> <td style="vertical-align: top;">1-43</td> </tr> </table>			Category ⁹	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³	X	FR, 2265-273 (LAB SOOTHERAP) 29 March 1974 See the translation	1-43	Y	FR, 2265-273 (LAB ZOOTHERAP) 29 March 1974 See the translation	1-43	Y	US, A, 4,444,777 (FLEMING, JR) 24 April 1984 See column 3, 4, and 5	1-43	Y	US, A, 4,537,894 (BLANCHARD) 27 August 1985 See column 4 lines 1-15.	1-43	Y	FR, 2390-959 (BRUGNAUD) 01 January 1979 See the translation.	1-43	Y	EP 0001-503 (NUNN) 18 April 1979 See page 2, lines 14-21	1-43
Category ⁹	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³																					
X	FR, 2265-273 (LAB SOOTHERAP) 29 March 1974 See the translation	1-43																					
Y	FR, 2265-273 (LAB ZOOTHERAP) 29 March 1974 See the translation	1-43																					
Y	US, A, 4,444,777 (FLEMING, JR) 24 April 1984 See column 3, 4, and 5	1-43																					
Y	US, A, 4,537,894 (BLANCHARD) 27 August 1985 See column 4 lines 1-15.	1-43																					
Y	FR, 2390-959 (BRUGNAUD) 01 January 1979 See the translation.	1-43																					
Y	EP 0001-503 (NUNN) 18 April 1979 See page 2, lines 14-21	1-43																					
<div style="display: flex; justify-content: space-between;"> <div style="width: 48%;"> <p>¹⁰ Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 48%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"A" document member of the same patent family</p> </div> </div>																							
IV. CERTIFICATION <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 50%; border-bottom: 1px solid black;">Date of the Actual Completion of the International Search</td> <td style="width: 50%; border-bottom: 1px solid black;">Date of Mailing of this International Search Report</td> </tr> <tr> <td style="border-bottom: 1px solid black;">17 DECEMBER 1990</td> <td style="border-bottom: 1px solid black; text-align: center; font-size: 1.2em;">11 FEB 1991</td> </tr> <tr> <td style="border-bottom: 1px solid black;">International Searching Authority</td> <td style="border-bottom: 1px solid black;">Signature of Authorized Officer <i>Nguyen</i></td> </tr> <tr> <td style="border-bottom: 1px solid black;">ISA/US</td> <td style="border-bottom: 1px solid black; text-align: center;">INTERNATIONAL DIVISION STANLEY J. FRIEDMAN</td> </tr> </table>			Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	17 DECEMBER 1990	11 FEB 1991	International Searching Authority	Signature of Authorized Officer <i>Nguyen</i>	ISA/US	INTERNATIONAL DIVISION STANLEY J. FRIEDMAN													
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report																						
17 DECEMBER 1990	11 FEB 1991																						
International Searching Authority	Signature of Authorized Officer <i>Nguyen</i>																						
ISA/US	INTERNATIONAL DIVISION STANLEY J. FRIEDMAN																						